



The American College of Obstetrics and Gynecology (ACOG), Society of Maternal Fetal Medicine (SMFM), American Society of Reproductive Medicine (ASRM) and CDC recommend that pregnant women, breast feeding women and women who are considering pregnancy receive the Covid vaccine.

The CDC and FDA have recommended a booster dose for people who have received the Pfizer Covid vaccine and are in a high-risk group. The full list of high-risk groups can be found on the CDC website, although includes people who are pregnant, post-partum and people who work in healthcare. The Pfizer booster is recommended 6 months after the 2nd vaccine was received. It is not known at this time if a booster is needed for people who have received the Moderna or Johnson & Johnson Covid vaccine.

At this time, someone is considered fully vaccinated 2 weeks following the second dose of Pfizer or Moderna vaccine or 2 weeks following the single dose of Johnson & Johnson Covid vaccine.

At Alliance, we support the use of Covid vaccine for all of our patients. We can discuss with you at your next appointment the use of vaccines in pregnancy and lactation as well as your risk of infection.

Alliance does not provide Covid vaccine exemption for people who are pregnant, breast feeding or trying for pregnancy.

CDC – Booster FAQ

<https://www.cdc.gov/coronavirus/2019-ncov/vaccines/booster-shot.html#FAQs>

Society for Maternal Fetal Medicine:

https://s3.amazonaws.com/cdn.smfm.org/media/3089/Provider_Considerations_for_Engaging_in_COVID_Vaccination_Considerations_8-23-21.pdf

American College of Obstetrics and Gynecology:

<https://www.acog.org/clinical/clinical-guidance/practice-advisory/articles/2020/12/covid-19-vaccination-considerations-for-obstetric-gynecologic-care>

American Society of Reproductive Medicine:

<https://www.asrm.org/news-and-publications/news-and-research/press-releases-and-bulletins/new-study-reveals-covid-vaccine-does-not-cause-female-sterility/>